Morbidity and Mortality

Weekly Report

PUBLIC HEALTH SERVICE

U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFAR

Prepared by the

COMMUNICABLE DISEASE CENTER

MElrose 4-5131

SEP 28 1961

CDC LIBRARY

For release September 29, 1961

Atlanta 22, Georgia

Vol. 10, No. 38

Provisional Information on Selected Notifiable Diseases in the United States and on Deaths in Selected Cities for Week Ended September 23, 1961

Poliomyelitis - Total and paralytic cases of poliomyelitis reported this week decreased by about one-third. A total of 71 cases, including 41 paralytic, were reported. This is in contrast to the 98 cases, 62 paralytic, reported during the previous week.

Both total and paralytic cases reported through the first 38 weeks are less than half the cases reported in the corresponding period last year. This is shown in the following table:

Poliomyelitis Cases (Cumulative) Through 38th Week

	1961	1960	1959	1958	1957
Paralytic	554	1446	3771	1680	1478
Total	849	2111	5950	3433	4658

New York has accounted for at least one-third of the total cases reported in the United States for the fourth time in the last five weeks. The tri-county outbreak in Upstate New York remains the only significant concentration of poliomyelitis in the United States. (See Epidemiological Reports).


Hepatitis – For the second consecutive week there was an increase in the number of reported hepatitis cases. For the 38th week 1213 cases were reported; 1130 cases had been reported the preceding week. It is too early to say whether or not this represents the beginning of the expected seasonal increase in cases.

Encephalitis - The number of cases of encephalitis remains at an unusually low level.

Table I. Cases of Specified Notifiable Diseases: United States

(Cumulative totals include revised and delayed reports through previous week)

Disease		38th Wee	k	The same		Cumulat	ive			OF ASSESSED
(Seventh Revision of International Lists, 1955)	Ended	Ended	2 P. St.	F	rst 38 wee	ks	Since s	Approxi- mate		
* Weekly incidence low or sporadic Data not available - Quantity zero	Sept. 23, 1961	Sept. 24, 1960	Median 1956-60	1961	1960	Median 1956-60	1960-61	1959-60	Median 1955-56 to 1959-60	seasonal low point
Anthrax062	2.421	2	*	5	15	*	*	*	*	*
Botulism049.1			*	5	10	*	*	*	*	*
Brucellosis (undulant fever)044	15	15	15	454	588	595	*	*	*	*
Diphtheria055	12	19	20	401	469	540	92	130	143	July 1
Encephalitis, infectious	38	58	76	1,173	1,389	1,426	1,173	1,389	1.426	Jan. 1
Repatitis, infectious, and	134934	707 -61	MA WOULD				, -,	-,		
Berum	1,213	898	260	56,164	27,810	14,825	3,343	2,269	806	Sept. 1
Malaria110-117	1	6	*	43	53	*	*	*	*	*
measles085	627	733	857	387,689	402,115	449,901	2,061	2,250	2,661	Sept. 1
Meningitis, esentic	141	166		2,143	2,046	100	2,143	2.046		Jan. 1
"eningococcal infections057	29	34	38	1,591	1,630	1,798	76	96	102	Sept. 1
Poliomyelitis080	71	203	431	849	2,111	4,650	746	1,901	4,122	Apr. 1
Paralytic080.0,080.1	41	134	218	554	1,446	1,471	494	1,289	1,196	Apr. 1
Nonparalytic080.2	22	44	141	200	460	2,421	175	433	2,260	Apr. 1
Unspecified	-8	25	72	95	205	758	77	179	666	Apr. 1
Falttacosis		-1	*	53	73	*	*	*	*	1 1 1
Streptococcal sore throat.	1 2 2 4		*	3	2	*	*	*	91 *	b resign
Including scarlet fever050.051	3,209	3,353		242,918	233,760	0.10	23,707		Indeed 1	Aug. 1
"VPROID fever	20	27	26	580	592	747	466	463	567	Apr. 1
Typhus fever, endemic101	2	2	*	29	53	*	*	*	*	*
Rabies in animals	61	46	74	2,570	2,733	3,337	3,146	3,701	4,279	Oct. 1



EPIDEMIOLOGICAL REPORTS

Diphtheria in the U. S.: Present Trends

The occurrence of diphtheria in the United States during the present year is closely following the patterns observed regularly in past years. Although the total of 401 cases reported to date during 1961 has declined slightly when compared with previous years, this decrease is not great.

The concentration of diphtheria in the southern States continues to be striking. The map depicts this distribution for diphtheria cases in 1960. During 1961 this same geographic distribution is apparent. Of 401 reported cases, 311 (77.6%) have occurred in the 17 states comprising the South Atlantic, East South Central, and West South Central Divisions of the country. The sole consistent exception to the low level of diphtheria case incidence in the other states is Minnesota where during the past several years diphtheria associated with infection by a gravis strain C. diphtheriae has been repeatedly observed. A total of 24 cases has occurred in Minnesota so far this year.

The coming of autumn in past years has regularly been accompanied by a striking increase in the occurrence of diphtheria. This is demonstrated in the graph in which cases during 1959 and 1960 are charted by month of onset. It is expected that this seasonal pattern will be following during 1961 to the same striking degree, although the rise in cases has not yet been reflected in the weekly state morbidity reports.

Preliminary observations of diphtheria cases reported to Surveillance Section, CDC, with onsets during 1961 reaffirms the fact that the great bulk of cases, particularly severe and fatal cases, is occurring in unimmunized children (See MMWR Vol. 10, No. 25). This is shown for 1960 cases in the following table:

Immunization Status	Cases	Percent	Deaths	Percent
Complete	55	10	0	0
Lapsed	46	8	0	0
Inadequate	56	10	3	8
None	409	72	37	92
Total Confirmed Cases with information on	James El	cment to	100 Hadi 100 Hadi	a process
Immunization Status	566	100%	40	100%

Poliomyelitis - New York State

Through Monday, September 25, 71 cases of paralytic poliomyelitis have occurred in the contiguous counties of Onondaga, Madison, and Oneida in Upstate New York. Of the 71 cases, 41 have occurred in Onondaga County (29 in Syracuse), 17 in Madison County, and 13 in Oneida County. The 67 paralytic cases with onset of symptoms since August are shown by county:

Week Ending	8/5	8/12	8/19	8/26	9/2	9/9	9/16	9/23
Onondaga County	2	1	4	11	11	3	6	12
Madison County	4	3	5	2	dil Inc	1	0	0
Oneida County	2	2	2	3	1	0	2	0

In addition to the 71 paralytic cases, 22 nonparalytic cases have been recognized in the tri-county area.

(Continued on page 8)

Table 2. CASES OF SPECIFIED NOTIFIABLE DISEASES: UNITED STATES, EACH DIVISION AND STATE, AND PUERTO RICO, FOR WEEKS ENDED SEPTEMBER 24, 1960 AND SEPTEMBER 23, 1961

(By place of occurrence. Numbers under diseases are category numbers of the Seventh Revision of the International Lists, 1955)

or heart 1975				Po	liomyeli	tis 080	(1) 10 (Eq.)				Menin-	Brucel losis
	(Includ		otal ot specifie	d by type)	Par	alytic (080.0,080	0.1	Nonpar	alytic	gitis, aseptic	(undu-
Area	38th	Week		Cumulative, first 38 weeks		38th Week		lative, 38 weeks	080	50.0 4 .00000	340 pt.	fever)
(20) (3) (4) (4)	1961	1960	1961	1960	1961	1960	1961	1960	1961	1960	1961	1961
UNITED STATES	71	203	849	2,111	41	134	554	1,446	22	44	141	15
NEW ENGLAND	1	14	18	186	1	14	15	147		- 54	11	
Maine		9	1	30	-)	9	1	30	12-		2	100
New Hampshire	-	-	1	7.3	- 1	1 2	-		-	-	-	-
Vermont	1	1 -	3	6		1	3	2	-	-	1	-
Rhode Island	_	- []	-	26 96	1	17-	8	18	3	-	4	-
Connecticut		4	4	28	2	4	3	74 23	-		5	
	- 00			2			1 1				-	
IDDLE ATLANTIC	30	35	215	292	17	18	150	213	10	17	8	-
New York	24 3	22	155 31	170	13	10	101	114	8	12	1	-
Pennsylvania	3	9	29	61 61	1 3	2 6	26	46	2	2	7	-
to be comed because of the state of the state of		LOT N	11 11 25	9			23	53	15.	3	7	-
AST NORTH CENTRAL	10	41	94	352	7	26	56	197	2	10	46	2
Ohio	3	13	25	86	1	6	12	38	1	3	5	1
IndianaIllinois	1	12	11 21	75 111	1	7	6	54	-	1	2	
Michigan	1	3	18	58	1	10	9	74	1	2	13	1000
Wisconsin	4	4	19	22	4	1	15 14	26 5	7.	1	24	1
AND DESCRIPTION OF THE PROPERTY OF THE PROPERT		25		2.0						3	2	-
Minnesota	9 1	11	52	120 36	4	5	24	63	4	6	34	7
Iowa	5	5	6 18	19	1 3	1	6 9	29			23	
Missouri	1	4	12	24	-	3	3	4	1	4	6	3
North Dakota		8 1 -	2	9	100		2	15 3	1	1	3	
South Dakota		6	1 i	4	120		345	1			1 2000	1
Nebraska	2	1	6	12		F_	3	7	2	1		-
Kansas	-	-	7	16	_	-	3	4		-	2	3
SOUTH ATLANTIC	5	33	142	342	2	30	103	253	3	3	9	2
Delaware	-	-	2			-	1	255			9	2
Maryland	1 1 40	17	23	51	-	16	23	45	-1-	1	100	ale al
District of Columbia			1	1 -1	1.5	- I	1	17.			100 702	-
Virginia	1	2	10	17	1	2	8	15	100		4	2
West Virginia	3	3	21	32	1	2	13	25	2	1	5	
North Carolina	1	2	14	67	100	1	7	47	1	1		-
South Carolina		6	15	110	1	6	10	73	7 7	-	100	
Georgia	A 1 14	1 2	27	15	1400	1	20	13	3	-		18 m
Florida			29	50		2	20	35	15.	-		-
AST SOUTH CENTRAL	4	30	64	146	2	9	39	68	1	1	3	
Kentucky	-	20	21	76			5	5		-	100	-
Tennessee	3	7	16	29	1	6	8	22	1	1	3	
Alabama	1	2	9 18	13 28	1	2	9	13				-
Agreement agreement hands to be built					0.00	1	17	28	10.0	1	The state of	-
EST SOUTH CENTRAL	5	12	114	208	3	9	60	125	2	3	3	3
Arkansas	4	3	15	21	2	3	6	13	2	-	1	2
Louisiana	- 100	4	36 3	42	50	3	27	27		1	-	
Oklahoma	- 1	I 5	60	12 133	1202	3	27	8	-	-	-	1
Philippin and the second of th			OT 9	973	The second		100	77	-	2	2	
OUNTAIN	1	3	41	60	-	3	23	27	-	-		
Montana		0 1	3	14	301	1	2	10	120	-	-	
Wyoming	1	-	14	5 18	0+)		6	1	-	-		-tagili
Colorado	or Harr	2	6	18	1/21	2	6	10	- 15-	4445	L. Wild	وفيده
New Mexico	110		3	4	112-	-		10	1	Section 1		0.00
Arizona	1.8	2 1 1	7	4	1.25	- 14	5	- 4		(in the last of the	11-11-12	ISL/NA
Utah	1.55	1.4	8	4		1 7	4	1	3000	200	-	-4610
Nevada	100	J- 1	P	100	1.2	100	-	132.1	-	-		pources.
ACIFIC	6	24	109	405	5	20	84	353	1	4	27	1
Washington	1	5	20	27		5	14	27			5	اخلول
Oregon	1	2	13	28	1	1	6	16	1-11-	1	1	1250
California	4	17	73	341	4	14	61	301		3	21	1
Alaska	-5	1 - 1	-	2	1.5	-	-	2	- 1	-	1	-14
Hawaii		0 1	3	7	1870	-	3	7	-			Tales -
Section 1		2			100		1					- 1
uerto Rico	-	16	6	448	-011	16	6	441	-	-		45.14

Table 2. CASES OF SPECIFIED NOTIFIABLE DISEASES: UNITED STATES, EACH DIVISION AND STATE, AND PUERTO RICO, FOR WEEKS ENDED SEPTEMBER 24, 1960 AND SEPTEMBER 23, 1961 - Continued

(By place of occurrence. Numbers under diseases are category numbers of the Seventh Revision of the International Lists, 1955)


-terril		Diphthe	eria 055			alitis, tious		titis, i erum 092			Meas	les
Area	38th	Week	Cumulative, first 38 weeks		082		38th Week		Cumulative, first 38 weeks		085	
	1961	1960	1961	1960	1961	1960	1961	1960	1961	1960	1961	1960
UNITED STATES	12	19	401	469	38	58	1,213	898	56,164	27,810	627	733
NEW ENGLAND	4	d.	6	10	4	1	78	14	1,703	832	27	131
Maine	-	T. 1-	-	2	-	-	5	1	109	50	13	2
New Hampshire	-	-	-	-	-	- 1	12	-	152	25	5	6
Vermont	3.5	-		-		- 1	8	1	174	12		33
Rhode Island	L L		5 -	7 1	1 1	1	32 4	7	668	414		80
Connecticut			1	_	2	-	17	1 4	205 395	162 169	9	10
	_	_								1		
MIDDLE ATLANTICNew York	118		20	13 3	4	6	156	191	7,724	3,296	75	76
New Jersey	1 1000	1 -	' '	2		4 2	71 27	129	3,273 1,850	1,785 222	44 13	45 9
Pennsylvania		1 1 2	13	8	4	_	58	53	2,601	1,289	18	22
EAST NORTH CENTRAL	1		15	36	5	17	201					
Ohio	CH	4	1	15	5	5	76	145	11,227 3,776	5,042 1,714	162 20	148 19
Indiana	_		1	5		11	15	15	1,715	568	26	17
Illinois	-	-	10	5			43	38	1,970	1,061	35	10
Michigan	41 30	- Lui-	3	9	-	1	64	28	3,489	1,520	43	36
Wisconsin	-	-	- 1	2	-	-	3	5	277	179	38	66
WEST NORTH CENTRAL		1	35	25	11	5	86	53	5,443	1,949	12	31
Minnesota	_	1	24	7		_	19	14	1,186	230	- 12	, ,,,
Iowa	-	-	1	7	-	3	32	13	1,594	332	3	8
Missouri	-	-	- 1	2	1	1	23	11	1,226	709	7	1
North Dakota		-	2	1			- 0	- I	126	141	1	21
South Dakota	-	-	6	5	-	1	6	2	147	129	1	-
Nebraska	- 1° l	-	2	1	2	-	6	8	561	207	-	1
Kansas		-		2	8	-1-4	- I	5	603	201	NN	NN
SOUTH ATIANTIC	8	7	88	140	5	11	155	80	6,987	3,253	69	59
Delaware	1	101-	-		1	-	8	6	166	200	-	2
Maryland			1	1	1	. 9	15	14	639	336	29	3
District of Columbia		_	2	10	-		6	1 :	89	40		
Virginia	10		15 1	18 4	1 711	1	16 32	5 20	1,103	634	10	13
West Virginia	m ver	1 4	7	. 7			43	8	1,311	611 281	19 1	19 1
South Carolina	3	1	8	43		1	7	1	358	51	1	6
Georgia	1	ī	23	22			6	4	624	214	-	-
Florida	4	3	31	45	3	_	22	22	1,173	886	9	15
EAST SOUTH CENTRAL		1	32	47	2	5	214					
Kentucky		1	8	1	2	1	55	143 57	8,438 2,479	3,992 1,472	18	24
Tennessee			3	7		3	85	43	3,345	1,297	14	20
Alabama		1	16	22	4	1	29	21	1,468	863	1	20
Mississippi	-		5	17	2	-	45	22	1,146	360	ī	_
WEST SOUTH CENTRAL	4	10	191	162	6	5	90	45	4,089	2,235	79	69
Arkansas		6	4	11	i	1	23	2	810	115	72	1
Louisiana	XIII.	2	23	33	-		-	6	425	118	214-0-1-0-2	1
Oklahoma	2	2	7	16	-31	- 0	6	6	275	280	-	-
Texas	2	110-	157	102	5	4	61	31	2,579	1,722	79	67
MOUNTAIN	_	-	8	35	-	3	56	52	3,385	2,241	64	57
Montana		_	2	3	_		6	6	304	103	24	4
Idaho	-	-	-	11	-2	- 1	8	2	249	262		3
Wyoming	1.6-1	-	-	5	-	2	1	-	136	23	-	-
Colorado	- tau 11 T	-	4	3	-	1	18	15	1,141	814	7	16
New Mexico	-		1	4	-	1.54	6	7	371	270	NN	
Arizona	N the re	11 6	14 15	3	-	1	3	11	520	496	19	15
UtahNevada	100	e lod	1	6			5	5	545	198	14	17
			F		1 7		9	6	119	75	-	
PACIFIC	H Can	-	6	1	1	5	177	175	7,168	4,970	121	138
Washington	4 1 5 5	-			1 -11	2	22	24	812	563	13	27
OregonCalifornia		3-	-			-	26	32	1,091	813	26	44
Alaska		1	2 /	-	1	3	117	114	4,881	3,367	70	63
Hawaii	4-4-	44	4	I .		11.24	12	3	333 51	153 74	12	1
Puerto Rico			48	110			10				70	,
	1,100	- 8	40	110	-	1	10	6	742	597	76	3

NN-Not Notifiable

Table 2. CASES OF SPECIFIED NOTIFIABLE DISEASES: UNITED STATES, EACH DIVISION AND STATE, AND PUERTO RICO, FOR WEEKS ENDED SEPTEMBER 24, 1960 AND SEPTEMBER 23, 1961 - Continued

(By place of occurrence. Numbers under diseases are category numbers of the Seventh Revision of the International Lists, 1955)

	Malaria	Meningo infec	occocal ctions	Psitta- cosis	Strepto- coccal sore throat,	I	yphoid :	fever 040)	Typhus fever, endemic		es in
Area	110-117	05	57	096.2	etc. 050,051	38th	Week		ative, 88 weeks	101	anı	MALIS
	1961	1961	1960	1961	1961	1961	1960	1961	1960	1961	1961	1960
UNITED STATES	1	29	34		3,209	20	27	580	592	2	61	46
NEW ENGLAND		2			102	-,	7	10			hon	3
Maine			1 2	-	103	1	-	18	8 2		-	
New Hampshire	_	_	_		1	1.00		-				1 -
Vermont	-	-	-	100	100	-	-	-	-	-	_	-
Massachusetts	1 -	/1/		100	36	1	-	12	3	-	1000	-
Rhode Island	1	- ·	7-327		5	-	-	2	-	-	-	-
Connecticut	-	1	- V	- P	61	-	-	3	3	- 10	-	-
MIDDLE ATLANTIC	1	5	5	-	90	2	1	70	42	-0	4	6
New York	1	1	4	-	53	1	1	35	28	- 1	4	6
New Jersey		1	1	-	16	-	-	17	1	- 79	-	-
Pennsylvania	- 1	3	-	-	21	1	-	18	13	-	-	-
EAST NORTH CENTRAL		14	7	-	184	3	1	78	76	-	4	4
Ohio	-		- 1	-	21	1	1	26	20	-	3	-
Indiana	1			-	80	- 1	-	19	22		1000	1
Illinois	After 1	6	1		14	W0.14	OHE N	25	20	-	1	-
Michigan	-	7	5		36	1	-	5	9	-	-	3
Wisconsin	-	1	1		33	1	-	3	5		-	-
WEST NORTH CENTRAL	-	1	4	-	110	2	2	31	36	-	14	9
Minnesota	- Day - 1			-	1		- C	5	1		4	_
Iowa	Name of	-	1	777.0	17		110000	1	6	ALCOUR TO	2	2
Missouri	2 252	1	50	-	35	2	2	19	21	to the end	5	5
North Dakota	NINO TO	V/ 5	4	DOTE SEE	18		5-1	11000	1	- T	1	1.5
South Dakota		13. E.				- N	-	3	3	-	2	2
Nebraska	-	300 211	3	-	100 -1	I SWIE DI	7 8 8 2	1	2	EXHIBIT S	11 14	
Kansas	100	100	119.00	51 F	39	444 1 T	10.00	2	2	German 16 T	0/160	1.000
SOUTH ATLANTIC	IN THESE	er futer	3	merks.ad	353	2	6	92	91	1	4	10
Delaware	- Sec. 6		Arm Land		1			1	1	- 1	T. 1	-
Maryland		100	1	(C)	8	-	1	2	4		-	-
District of Columbia	-	-	71 -11	-	2	(C)		10	7	agrumates.	30 30	100
Virginia	1000	C 385		179 H.	120 116	411	2 2	14	20	No. of	1	2
West Virginia		F 10-19	fermion a		10		_	9 13	9 8	1	1797	6
South Carolina		-	1		25	2		8	11	N. No. THE		
Georgia	7281107	3 11 1	1777	_	2	73 -0	1	25	21	Secretary 1	ST. Dice	1
Florida-	555 31-	1.52	1	-	69	W 1-	-	10	10	- I	3	1
EAST SOUTH CENTRAL	drantes.	9 1	2	1000	795	2	7	59	82		3	4
Kentucky	_	Sur Fan	1	_	36	2		12	15	1 - 2	1	3
Tennessee		1	-	1.0	746	-	6	39	48	200	2	1
Alabama	7.0000	1	BAST-1	II fren	5	a allero	i	6	14		4 1	_
Mississippi	-	3	1	-	8	1 11 -	-	2	5	-	-	-
WEST SOUTH CENTRAL	_	2	2	_	506			126	171	,	2,4	_
Arkansas		2			506	3	6 2	126 21	171 39	1	24	5
Louisiana	GET KI	DOMESTIC.	1	250 1907	3	CION W	2	16	54		10	2
Oklahoma	-	-	-	-	8	-	2	12	12		_	-
Texas	e many a b	-	1	A	495	1	2	77	66	1	14	3
MOUNTAIN			3		772	3		52		_		0
Montana	100	lary .	1		25	_		16	31 9		3	2
Idaho	4	100	15,55	Same	47	e. fe d		1	2			
Wyoming			1000		-	1 9-		3	4			
Colorado	0 (14)		2	-	329			6	mark.		1	_
New Mexico-	- 1	-		123 -	157	3	+ 7 -	14	8	-	1	2
Arizona	-		-00-	10035	107		-	6	7		2	-
Utah-		-	-		104		-	2	1	-	-	-
Nevada	-	1-1	-	-	3	-	-	4	-		-	-
PACIFIC	- J u	4	8	en, E.	296	1	4	54	55	Boltz 1	5	6
Washington	-	-			105	-	-	7	5	-		-
Oregon-		1	2	100 -0	21	and the	-	1	8	-	- 1	-
California-	-	3	5	= 1: J=	144	1	4	46	41	-	5	6
Alaska	-		1	10/47	19	100	-	-	1		-	-
Hawaii	7.0	-	-	237	7	44.07	14	-	.2.	-	-	-
D	100	176			Inigi.	MATELL	(2)					1111
Puerto Rico		li - 31	1 - 1	Con - 3	9	10000	11-1-	17	17	- Sec.		2



The chart shows the number of deaths reported for 117 major cities of the United States by week for the current year, a 5-week moving average of these figures plotted at the central week, and an adjusted average for comparison. For each region the adjusted average was computed as follows: From the total deaths reported each week for the years 1956-1960, 3 central figures were selected by eliminating the highest and lowest figure reported for that week. A 5-week moving average of the arithmetic mean of the 3 central figures was then computed with adjustment to allow for population growth in each region. The average value of the regional increases was 2 percent which was incorporated in the adjusted average shown in the chart.

Table 4 shows the number of death certificates re-

ceived during the week indicated for deaths that occurred in selected cities. Figures compiled in this way, by week of receipt, usually approximate closely the number of deaths occurring during the week. However, differences are to be expected because of variations in the interval between death and receipt of the certificate and because of incomplete reporting due to holidays or vacations. If a report is not received from a city in time to be included in the total for the current week, an estimate is used.

The number of deaths in cities of the same size may also differ because of variations in the age, race, and sex composition of the populations and because some cities are hospital centers serving the surrounding areas. Changes from year to year in the number of deaths may be due in part to population increases or decreases.

Table 3. DEATHS IN SELECTED CITIES BY GEOGRAPHIC DIVISIONS

(By place of occurrence and week of filing certificate. Excludes fetal deaths. Data exclude figures shown in parentheses in table 4)

	38th week	37th week ended Sept. 16, 1961	Adjusted average, 38th week 1956-60	Percent change, adjusted average to current week	Cumulative, first 38 weeks			
Area	ended Sept. 23, 1961				1961	1960	Percent change	
TOTAL, 117 REPORTING CITIES	10,776	11,035	10,634	+1.3	435,317	438,615	-0.8	
New England(14 cities)	611	706	634	-3.6	26,684	27,356	-2.5	
East North Central(21 cities)	2,993* 2,336*	3,124 2,384	2,902 2,295	+3.1	124,167 93,054	121,283 94,948	+2.4	
West North Central(9 cities) South Atlantic(11 cities)	748 883	768 884	740 883	+1.1	29,657 37,578	30,565 37,681	-3.0	
East South Central(8 cities)	505	565	491	+2.9	19,707	19,962	-1.3	
West South Central(13 cities) Mountain(8 cities)	1,034 361	907 360	971 341	+6.5	37,249 13,870	38,509 13,747	+0.9	
Pacific(13 cities)	1,305*	1,337	1,377	-5.2	53,351	54,564	-2.2	

^{*}Includes estimate for missing reports.

Table 4. DEATHS IN SELECTED CITIES

(By place of occurrence and week of filing certificate. Excludes fetal deaths)

denomination Area	38th week ended Sept.	37th week ended Sept.	Cumula first 38	tive, Weeks	Area	38th week ended Sept.	37th week ended Sept.	Cumula first 38	
	23, 1961	16, 1961	1961	1960	Seit-Hei	23, 1961	16, 1961	1961	1960
meeting Authorate Mary	10.57		2702		See 12 - 12 - 12 - 12 - 12 - 12 - 12 - 12	35 7.6	325000	e Artina	8
EW ENGLAND:	11.131		ESTRICTION	2.1810,274	WEST NORTH CENTRAL Con.:				and a
Boston, Mass.	221	264	9,162	9,498	St. Louis, Mo	214	226	8,946	9,444
Bridgeport, Conn	36	42	1,457	1,568	St. Paul, Minn.	69	60	2,503	2,665
Cambridge, Mass	25	36	1,104	1,192	Wichita, Kans.	54	48	1,757	1,735
Fall River, Mass	30	19	1,021	1,076					
Hartford, Conn	40	43	1,820	1,834	SOUTH ATLANTIC:	11.00	2.5		
Lowell, Mass	25	15	922	909	Atlanta, Ga	86	111	4,286	4,51
Lynn, Mass.	24	28	827	916	Baltimore, Md	213	214	9,355	9,58
New Bedford, Mass	18	26	986	935	Charlotte, N.C	27	33	1,347	1,48
New Haven, Conn	36	43	1,740	1,703	Jacksonville, Fla	50	50	2,171	2,27
Providence, R.I	45	63	2,362	2,425	Miami, Fla	78	59	2,889	2,78
Somerville, Mass	8	15	500	504	Norfolk, Va	44	41	1,892	1,53
Springfield, Mass	39	34	1,687	1,709	Richmond, Va	54	62	2,946	2,97
Waterbury, Conn	16	26	1,018	1,040	Savannah, Ga	38	31	1,240	1,29
Worcester, Mass	48	52	2,078	2,047	St. Petersburg, Fla	(58)	(56)	(2,600)	(2,71
Today Tolerout much	il negli	or well the	The Check	400	Tampa, Fla	58	48	2,530	2,49
IDDLE ATLANTIC:			1 700	1 (57	Washington, D.C.	196	195	7,408	7,29
Albany, N.Y	40	35	1,728	1,657	Wilmington, Del	39	40	1,514	1,44
Allentown, Pa	35	32	1,311	1,316		O THE			
Buffalo, N.Y.	138*	130	5,573	5,567	EAST SOUTH CENTRAL:	1 3500	3.63	THE REAL PROPERTY.	1016, 27
Camden, N.J.	36	47	1,625	1,618	Birmingham, Ala	90	77	3,246	3,23
Elizabeth, N.J.	33 26	42 34	1,119	1,120 1,479	Chattanooga, Tenn	51	49	1,783	1,79
Jersey City, N.J	54	63	2,635	2,701	Knoxville, Tenn	39	35	1,056	1,06
Newark, N.J.	82	75	3,826	3,679	Louisville, Ky	86	142	4,356	4,37
New York City, N.Y	1,484	1,625	63,304	61,577	Memphis, Tenn	100	104	4,299	4,29
Paterson, N.J.	29	40	1,481	1,455	Mobile, Ala Montgomery, Ala	43	45	1,537	1,58
Philadelphia, Pa	476	490	19,565	18,611	Nashville, Tenn	43	46	1,213	1,32
Pittsburgh, Pa	211	186	7,249	7,357	Mashville, Telli,	53	67	2,217	2,30
Reading, Pa	19	25	880	898	WEST SOUTH CENTRAL:	130 6H	Decky	18,772	
Rochester, N.Y	104	105	3,838	3,804	Austin, Tex	33	29	1,294	1,33
Schenectady, N.Y	27	16	911	909	Baton Rouge, La	30	27	1,039	1,09
Scranton, Pa	40	26	1,347	1,417	Corpus Christi, Tex	22	27	838	90
Syracuse, N.Y	60	65	2,328	2,344	Dallas, Tex	131	115	4,757	4,75
Trenton, N.J.	39	45	1,709	1,572	El Paso, Tex	42	29	1,333	1,46
Utica, N.Y	32	22	1,071	1,035	Fort Worth, Tex	58	77	2,432	2,55
Yonkers, N.Y	28	21	1,179	1,167	Houston, Tex.	222	128	6,273	6,44
		= V 1 -			Little Rock, Ark	64	69	2,156	2,20
AST NORTH CENTRAL:		12.3	W-5.6		New Orleans, La	161	153	6,394	6,86
Akron, Ohio	75	56	2,173	2,169	Oklahoma City, Okla	81	61	2,838	2,85
Canton, Ohio	31	32	1,169	1,322	San Antonio, Tex	99	99	3,885	3,85
Chicago, Ill.	728	659	28,175	29,339	Shreveport, La	39	46	1,927	2,07
Cincinnati, Ohio	140	160	5,933	5,966	Tulsa, Okla	52	47	2,083	2,10
Cleveland, Ohio	169	199	7,696	8,007	and said guidered of care	p Ruphy	to out	e judan	
Columbus, Ohio	110	129	4,346	4,483	MOUNTAIN:	1976	1,-1-	This Total	
Dayton, Ohio	84 308	78 328	3,071	2,832	Albuquerque, N. Mex	26	25	1,182	1,17
Detroit, Mich.			12,669	12,965	Colorado Springs, Colo	8	14	608	62
Evansville, Ind	36	39 48	1,379	1,381	Denver, Colo	116	120	4,356	4,51
First Wayne Ind	26 35	31	1,618	1,531 1,412	Ogden, Utah	18	14	637	63
Fort Wayne, Ind.	29	25	1,459	1,196	Phoenix, Ariz.	94	83	3,135	2,94
Grand Rapids, Mich.	33	53	1,710	1,572	Pueblo, Colo	17	15	638	63
	138	141	5,402	5,573	Salt Lake City, Utah	48	43	1,842	1,85
Madison, Wis.	33	34	1,250	1,207	Tucson, Ariz	34	46	1,472	1,36
Milwaukee, Wis	120	122	4,662	4,758	PACIFIC:				
Peoria, Ill.	28	29	1,072	1,150	Berkeley, Calif	71:10	Esca		S HE STATE
Rockford, Ill.	28*	25	1,072	1,091	Fresno, Calif	20	22	659	63
South Bend, Ind.	31	37	1,090	1,092	Glendale, Calif	(35)	(25)	(1,651)	(1,69
Toledo, Ohio	104	104	3,743	3,806	Honolulu, Hawaii	(30)	(34)	(1,262)	(1,47
Youngstown, Ohio	50	55	2,205	2,096	Long Beach, Calif	44	36	1,513	1,58
	30	1	_,	_,050	Los Angeles, Calif	44	44	2,075	2,09
EST NORTH CENTRAL:		and last		1 10	Oakland, Calif.	431	494	18,885	19,25
Des Moines, Iowa	58	58	2,042	2,089	Pasadena, Calif	89	93	3,664	3,66
Duluth, Minn.	21	36	987	960	Portland, Oreg	39	27	1,256	1,31
Kansas City, Kans	30	30	1,411	1,327	Sacramento, Calif	131	90	4,056	4,21
Kansas City, Mo	122	132	4,863	4,809	San Diego, Calif	38 93	60	2,355	2,21
Lincoln, Nebr	(25)	(32)	(1,051)	(990)	San Francisco, Calif	182	93	3,359	3,42
Minneapolis, Minn	108	108	4,513	4,731	San Jose, Calif		196	7,400	7,53
Omaha, Nebr	72	70	2,635	2,805	Seattle, Wash	108	(37) 108	(1,327)	(1,3
A STATE OF THE STA			1144		Spokane, Wash	43*		4,921	5,26
stimate - based on average	1224		7.5-		Tacoma, Wash	43*	48 26	1,762	1,79
								1,446	

Figures shown in parenthesis are from cities which have been reporting less than five years and hence are not included in Table 3.

Since the oral poliomyelitis immunization program began on August 29, there have been 25 cases of paralytic poliomyelitis. These cases have had onset as follows (3 day periods):

Orai	August	25-	September							
Vaccine Status	29-31*	1-3	4-6	7-9	10-12	13-15	16-18	Total		
No Vaccine	4	1	1	1	3	2	0	12		
Vaccine	1	6	1	1	0	3	1	13		

*Oral Vaccine administered during this three day period

Bat Rabies - Tennessee and Arkansas

Each of two States, Tennessee and Arkansas, recently reported its first laboratory confirmed case of rabies in bats.

On July 18, 1961, a university student's son found an injured bat under a tree at a Vanderbilt University graduate student housing facility. The bat was kept until it died on July 24, and then submitted to the State Department of Public Health for laboratory analyses. Examination for Negri bodies was negative but rabies was confirmed by fluorescent antibody technique.

Identification of the species of bat was made by the father and son by inspection of a collection of 14 species of indigenous bats. It was felt that this collection represented all species, except the hoary, reported in Tennessee. It was tentatively concluded that the rabid bat was a Nycteria borealis.

On August 17, 1961, a bat bit a woman resident of Little Rock, Arkansas. The bat had been lying enclosed in a small paper box in the front yard of this woman's residence. One of her children opened the box. The bat flew out and lit on the child's dress. In brushing the bat off the little girl's dress, the woman was bitten on the right index finger.

Examination of the bat's brain by fluorescent antibody technique at the Communicable Disease Center, Atlanta, Georgia made the diagnosis of rabies. The bat was not the dark type usually seen in Arkansas but was identified as an insectivorous bat which was light tan in color.

Several minutes after this episode was reported on a local television station, a man called to explain the mysterious presence of the bat in the paper box. He said that the bat in question had flown into his car on August 17. He had captured it in a paper box and discarded it. (Reported by Cecil B. Tucker, M.D., Director, Division of Preventable Diseases, Tennessee Department of Public Health, and Harvie R. Ellis, D.V.M., Director, Division of Veterinary Public Health, Arkansas State Health Department.)

SOURCE AND NATURE OF MORBIDITY DATA

These provisional data are based on reports to the Public Health Service from the health departments of each State and Puerto Rico. They give the total number of cases of certain communicable diseases reported during the week usually ended the preceding Saturday. Total figures for the United States and the Pacific Division include data for the States of Alaska and Hawaii. Cases of anthrax, botulism, and rabies in man are not shown in table 2, but a footnote to table 1 shows the States reporting these diseases. When diseases of rare occurrence are reported by a State (cholera, dengue, plague, louse-borne relapsing fever, smallpox, louse-borne epidemic typhus, and yellow fever) this is noted below table 1.

QUARANTINE MEASURES Immunization Information for International Travel

No Changes Reported

Library 81 CD0 7 61

PUBLIC HEALTH SERVICE
Communicable Disease Center
Atlanta 22, Georgia
Official Business

MAIL